

# Implementation Science: routine PROMs in cancer

November 2023



A good argument can be made that the evidence base on routine PROMs in some cancers is sufficiently mature to recommend routine deployment

For example...

Effectiveness of routine provision of feedback from patient-reported outcome measurements for cancer care improvement: a systematic review and metaanalysis

Sheng-Chieh Lu<sup>1</sup>, I Porter<sup>2</sup>, J M Valderas<sup>3 4 5</sup>, C J Harrison<sup>6</sup>, Chris Sidey-Gibbons<sup>7</sup>

**Results:** We included 29 studies involving 7071 cancer participants. A small number of studies was available for each metanalysis (median = 3 studies, ranging from 2 to 9 studies) due to heterogeneity in the evaluation of the trials. We found that the intervention improved HRQL (Cohen's d = 0.23, 95% CI 0.11-0.34), mental functioning (Cohen's d = 0.14, 95% CI 0.02-0.26), patient-healthcare provider communication (Cohen's d = 0.41, 95% CI 0.20-0.62), and 1-year overall survival (OR = 0.64, 95% CI 0.48-0.86). The risk of bias across studies was considerable in the domains of allocation concealment, blinding, and intervention contamination.

**Digi**Core

# A good argument can be made that the evidence base on routine PROMs in some cancers is sufficiently mature to recommend routine deployment

#### For example...

Effectiveness of routine provision of feedback from patient-reported outcome measurements for cancer care improvement: a systematic review and metaanalysis

Sheng-Chieh Lu<sup>1</sup>, I Porter<sup>2</sup>, J M Valderas<sup>3 4 5</sup>, C J Harrison<sup>6</sup>, Chris Sidey-Gibbons<sup>7</sup>

**Results:** We included 29 studies involving 7071 cancer participants. A small number of studies was available for each metanalysis (median = 3 studies, ranging from 2 to 9 studies) due to heterogeneity in the evaluation of the trials. We found that the intervention improved HRQL (Cohen's d = 0.23, 95% CI 0.11-0.34), mental functioning (Cohen's d = 0.14, 95% CI 0.02-0.26), patient-healthcare provider communication (Cohen's d = 0.41, 95% CI 0.20-0.62), and 1-year overall survival (OR = 0.64, 95% CI 0.48-0.86). The risk of bias across studies was considerable in the domains of allocation concealment, blinding, and intervention contamination.

Implementation Science

So how can we do this?

What barriers to solve?

What are the benefits?

How to reuse for research?



### Typical challenges to routine implementation and research reuse

Implementation challenges

- Funding: monitoring nurses, technology integration
- Workflow integration: care pathways, managing recall capacity
- Patient survey fatigue: simple & easy instruments vs complex, more accurate
- Multidisciplinary implementation: IT care pathway design and clinical

Research interoperability challenges

- Need: same instrument, same cohort, same point in pathway
- **Reality:** many instruments, highly confounded by timing in pathway



## **Our panel today**

Medizinische Universitat Innsbruck



Bernhard Holzner

The benefits and challenges of introducing PROMs into routine cancer care Cliniques Universitaires Saint-Luc



Cédric Van Marcke

St. Luc's experience introducing ICHOM standards in breast cancer Ludwig Maximilian University Hospital



**Karin Berger** 

Large scale PROM pilots in Germany

